



Rapti academy of health science guidelines on retrograde intra renal surgery

Santosh Kumar Jha

Assistant Professor, Department of Urology, Rapti Academy of Health Sciences, Dang, Nepal

Abstract

This study aims to establish comprehensive guidelines for the safe and effective performance of Retrograde Intra Renal Surgery (RIRS) at Rapti Academy of Health Sciences, with a focus on treating renal stones. The guidelines were developed through a systematic review of literature from 2000 to 2023, utilizing databases like PubMed, Embase, and Cochrane Library. A panel of expert urologists reviewed the evidence, covering areas such as patient selection, preoperative evaluation, surgical techniques, and postoperative care. Emphasis was placed on improving stone-free rates (SFRs), with rates reaching up to 90% in specific cases, while addressing complications such as ureteral perforation and postoperative fever. The guidelines provide detailed recommendations for minimizing risks and enhancing outcomes through standardized practices. By following these guidelines, urologists can optimize patient outcomes and reduce complications associated with RIRS, and ongoing advancements in technology will require periodic updates to the protocols.

Keywords: Retrograde intra renal surgery, renal stones, ureteroscopy, patient outcomes, guidelines, stone-free rate

Introduction

Retrograde Intra Renal Surgery (RIRS) has revolutionized the treatment of renal stones and other upper urinary tract pathologies. This minimally invasive technique offers several advantages over traditional open surgery and percutaneous nephrolithotomy (PCNL), including shorter hospital stays, reduced morbidity, and faster recovery times. As the popularity of RIRS continues to grow, there is a need for standardized guidelines to ensure optimal patient outcomes and minimize complications.

The Rapti Academy of Health Science has developed these comprehensive guidelines to address the various aspects of RIRS, from patient selection to postoperative care. These guidelines are intended to serve as a reference for urologists, both experienced and those new to the technique, to promote best practices in RIRS.

1. Overview of retrograde intra renal surgery (RIRS)

Retrograde Intra Renal Surgery (RIRS) has become a transformative procedure in the management of renal stones and other upper urinary tract pathologies. As a minimally invasive technique, RIRS provides several advantages over traditional methods such as open surgery and percutaneous nephrolithotomy (PCNL), including shorter hospital stays, reduced morbidity, and faster recovery times (Geraghty *et al.*, 2023)^[9]. Over the last two decades, the development of flexible ureteroscopy (FURS), coupled with advancements in laser lithotripsy and digital imaging technologies, has enhanced the safety and efficacy of RIRS, making it the preferred method for managing small to medium-sized renal stones (Kim *et al.*, 2020)^[11]. According to the European Association of Urology, RIRS is recommended as a first-line treatment for renal stones under 2 cm, especially for patients who prefer a less invasive option (Kim *et al.*, 2023)^[12].

2. Success rates and advantages of RIRS

The primary measure of success in RIRS is the stone-free rate (SFR), which can reach up to 90% depending on factors such as stone size, location, and patient anatomy (Kim *et al.*, 2020)^[11]. For stones located in the lower pole or proximal

ureter, RIRS has demonstrated excellent outcomes, often outperforming extracorporeal shock wave lithotripsy (ESWL) and achieving results comparable to PCNL but with fewer complications and a quicker recovery (Fankhauser *et al.*, 2021; Juliebø-Jones *et al.*, 2023)^[7, 10]. However, variability in outcomes can occur in more complex cases, such as those involving larger stones or challenging renal anatomy, where access and visualization may be limited (Şeçkiner *et al.*, 2020)^[16].

3. Complications and safety considerations

While RIRS offers several advantages, it also presents challenges. The use of ureteral access sheaths, which facilitate smoother procedures and faster stone retrieval, can increase the risk of ureteral injuries, especially in complex cases (Amorim *et al.*, 2024; Cooper *et al.*, 2020)^[1, 4]. Complications such as fever, infection, and bleeding remain common concerns, particularly when intrarenal pressure is not managed properly. Studies have shown that maintaining low-pressure irrigation during the procedure is critical for reducing the risk of severe complications, such as pyelovenous backflow, renal pelvis rupture, or sepsis (Asutay *et al.*, 2022; Lildal *et al.*, 2022)^[2, 15]. Proper management of irrigation and pressure is thus an essential aspect of ensuring patient safety during RIRS (Feng *et al.*, 2020)^[8].

4. Technological advancements in RIRS

In recent years, technological innovations have played a significant role in improving the outcomes of RIRS. The introduction of digital ureteroscopes has enhanced visualization during the procedure, allowing for more precise stone fragmentation and higher SFRs, particularly in difficult-to-reach areas of the kidney (Ventimiglia *et al.*, 2022; Juliebø-Jones *et al.*, 2023)^[10, 18]. The adoption of single-use ureteroscopes has further revolutionized the practice by minimizing the risk of cross-contamination and reducing maintenance costs, particularly in high-volume centers (Ventimiglia *et al.*, 2022)^[18]. These advancements are indicative of the ongoing evolution in RIRS, aimed at improving procedural efficiency and patient outcomes.

5. Guideline development for standardized practice

Recognizing the need for standardized practices, the Rapti Academy of Health Sciences has developed comprehensive guidelines for RIRS. These guidelines are based on an extensive review of current literature and expert consensus, covering patient selection, preoperative evaluation, surgical techniques, and postoperative management. By adhering to these guidelines, urologists can minimize procedural risks and optimize patient outcomes, ensuring a standardized approach to RIRS (Dasgupta *et al.*, 2021)^[5]. The guidelines reflect a global shift towards less invasive urological procedures and emphasize the importance of personalized patient care, as outlined by leading urological associations (Kim *et al.*, 2023)^[12].

As RIRS continues to evolve with ongoing technological advancements, the guidelines provided by the Rapti Academy will require regular updates to incorporate new innovations. Future improvements in flexible ureteroscopy design, stone fragmentation techniques, and pressure management systems will further enhance the safety and efficacy of RIRS. These guidelines serve as a critical reference for urologists in ensuring the best possible outcomes for patients undergoing RIRS (Denstedt *et al.*, 2024)^[6].

Methods and materials

1. Study design

This study was based on a comprehensive review of the literature, focusing on the development of standardized guidelines for Retrograde Intra Renal Surgery (RIRS). The guidelines were created through a systematic analysis of peer-reviewed articles, expert consensus, and clinical data from 2000 to 2023. The methodology involved reviewing existing surgical practices, patient outcomes, and complications associated with RIRS, with the goal of creating evidence-based recommendations for urologists.

2. Literature search and data sources

The data collection process involved a systematic search of electronic databases, including PubMed, Embase, and Cochrane Library. The search terms used included "retrograde intrarenal surgery," "flexible ureteroscopy," "renal stones," "upper urinary tract," and related keywords. The literature review focused on randomized controlled trials (RCTs), systematic reviews, meta-analyses, and high-quality observational studies relevant to RIRS. Articles were selected based on their relevance to patient selection criteria, preoperative evaluation, surgical techniques, and postoperative care in the context of RIRS.

3. Inclusion and exclusion criteria

Only studies that provided significant clinical evidence on RIRS, flexible ureteroscopy, and the treatment of renal stones were included. Studies that focused on unrelated surgical procedures or lacked comprehensive clinical data on patient outcomes, such as stone-free rates (SFR) and postoperative complications, were excluded. Additionally, non-English language articles and abstracts were excluded to ensure consistency and clarity in the interpretation of results.

4. Expert panel and consensus

A panel of experienced urologists specializing in endourology and RIRS was convened to review and analyze

the data collected from the literature. The panel comprised 10 experts from various academic and clinical institutions, all of whom had performed a minimum of 100 RIRS procedures. The experts used the Oxford Centre for Evidence-Based Medicine criteria to grade the level of evidence and assess the strength of the recommendations. Consensus was achieved through multiple rounds of discussions and iterative revisions of the guidelines.

5. Development of guidelines

The development of the guidelines followed a structured approach, focusing on key aspects of RIRS, including patient selection, preoperative evaluation, surgical techniques, and postoperative management. The guidelines aimed to address various aspects of RIRS, such as:

- **Patient selection:** Criteria for choosing suitable candidates for RIRS based on stone size, location, and comorbidities.
- **Preoperative evaluation:** Recommendations for imaging studies, urinalysis, and blood work to ensure patient safety and optimize outcomes.
- **Surgical technique:** Detailed procedural steps, including access, stone fragmentation, and retrieval techniques using flexible ureteroscopy and laser lithotripsy.
- **Postoperative care:** Best practices for monitoring, stent placement, and managing potential complications such as infection, bleeding, and ureteral injury.

6. Grading of evidence

Each recommendation was assigned a grade based on the level of evidence supporting it. The grading system, following the Oxford Centre for Evidence-Based Medicine, ranged from Level 1 (highest quality evidence, such as RCTs and meta-analyses) to Level 5 (expert opinion without explicit critical appraisal). This grading system ensured that the guidelines were based on the best available evidence while allowing for clinical judgment in areas where data were limited.

7. Statistical analysis

The findings from the literature were synthesized, and statistical analyses were performed where applicable to determine the stone-free rates (SFR), complication rates, and the efficacy of different surgical techniques. Data were compared across studies to identify trends and areas requiring further investigation. The expert panel reviewed the statistical findings to ensure their clinical relevance and applicability in developing practical guidelines.

Results

The review of the literature and expert panel consensus resulted in the development of comprehensive guidelines for Retrograde Intra Renal Surgery (RIRS). These guidelines aim to standardize the procedure and ensure optimal patient outcomes, covering key aspects such as patient selection, preoperative evaluation, surgical techniques, and postoperative care.

1. Patient selection and preoperative evaluation

The guidelines emphasize that RIRS is most suitable for treating renal stones smaller than 2 cm, particularly for stones located in the lower pole or proximal ureter. In cases of anatomical abnormalities or patients with bleeding

diathesis, RIRS is recommended due to its minimally invasive nature. However, contraindications such as active urinary tract infections, untreated coagulopathy, and large stone burdens (>2 cm) were outlined, requiring alternative approaches like percutaneous nephrolithotomy (PCNL). The guidelines stress the importance of thorough preoperative evaluation, including medical history, urinalysis, and imaging techniques such as non-contrast CT scans to assess stone characteristics and renal anatomy.

Table 1: Indications for RIRS in Renal Stone Management

Indication	Stone Size	Location
Primary treatment	<2 cm	Any
Lower pole stones	<1 cm	Lower pole
Bleeding diathesis	Any	Any
Anatomical abnormalities	Any	Any
Residual stones after other treatments	Any	Any

2. Surgical technique and learning curve

The surgical steps of RIRS include accessing the ureter, localizing the stone, fragmenting it with laser lithotripsy, and retrieving it. The use of flexible ureteroscopes combined with ureteral access sheaths was found to improve procedural efficiency and minimize operative time. The success of the procedure is measured by the stone-free rate (SFR), which was reported to be as high as 90% in cases involving small stones (<2 cm). However, variability in SFR was observed in more complex cases, such as larger stones or difficult anatomical locations.

The learning curve for RIRS is significant, as the procedure requires specialized skills. Surgeons typically need to perform at least 50 supervised RIRS procedures before achieving independent proficiency. Figure 1 illustrates the gradual improvement in surgical outcomes as a surgeon's experience increases, emphasizing the importance of extensive training in RIRS.




Fig 1: Learning curve for RIRS

The learning curve demonstrates that increased experience in performing RIRS procedures leads to improved stone-free rates, reduced complications, and shorter operative times. Surgeons with more than 50 completed cases showed higher proficiency, indicating that RIRS requires a substantial amount of hands-on practice to master.

3. Postoperative care and complications

Postoperative care guidelines recommend regular monitoring of the patient, with emphasis on pain management, infection prevention, and stent placement

when necessary. Ureteral stents are particularly useful in cases where ureteral dilation is significant, operative times are extended, or residual stone fragments remain. Discharge criteria include stable vital signs, absence of infection, and manageable pain levels. Follow-up imaging is recommended to confirm stone clearance.

A range of potential complications was identified during RIRS, including urinary tract infections (UTIs), bleeding, ureteral injury, and postoperative fever. These complications are addressed with specific management strategies, as detailed in Table 2: Common Complications of RIRS and Their Management.

Table 2: Common complications of RIRS and their management

Complication	Incidence (%)	Management
Ureteral perforation	1-4	Stent placement or surgical repair
Bleeding	2-4	Conservative management or angioembolization
Stone migration	3-15	Anti-retropulsion devices or alternative procedure
Fever/sepsis	2-7	Antibiotics, possible decompression
Ureteral stricture	0.5-1	Endoscopic dilation or surgical repair

These findings emphasize the need for a careful approach during surgery and diligent postoperative care to minimize complications.

Discussion

1. Clinical implications of RIRS

The guidelines developed for RIRS represent a significant step toward standardizing a procedure that has become the preferred choice for treating small to medium-sized renal stones. The high stone-free rates, particularly for stones smaller than 2 cm, confirm that RIRS is highly effective when patient selection and surgical techniques are properly managed (Kim *et al.*, 2023) [12]. The guidelines stress the importance of comprehensive preoperative evaluations, including imaging and patient history, to ensure successful outcomes.

However, the complexity of the procedure requires surgeons to acquire significant experience before achieving optimal results. As demonstrated in Figure 1, the learning curve for RIRS is steep, and extensive hands-on training is required for surgeons to gain proficiency. This highlights the need for ongoing education and support for urologists to ensure that they can perform RIRS safely and effectively.

2. Challenges in RIRS practice

While RIRS offers several advantages, it also presents challenges. One of the most significant is the use of ureteral access sheaths, which, although they facilitate stone retrieval, can increase the risk of ureteral injuries. As shown in Table 2, ureteral perforation is a rare but significant complication, and surgeons must be vigilant in ensuring proper insertion techniques and pressure management to avoid trauma (Cooper *et al.*, 2020; Amorim *et al.*, 2024) [1, 4]. Managing intrarenal pressure during the procedure is also critical, as high pressures can lead to serious complications like pyelovenous backflow or sepsis (Asutay *et al.*, 2022) [2]. Postoperative complications, such as fever and infection, are not uncommon, but they can be effectively managed

through proper stent placement and postoperative monitoring. As outlined in Table 2, postoperative infections (UTIs and sepsis) require prompt antibiotic treatment and, in some cases, decompression of the renal system.

3. Technological advancements and future directions

Technological advancements have played a key role in improving RIRS outcomes. The use of digital ureteroscopes provides superior visualization, enabling more precise stone localization and fragmentation. These advances, along with the development of single-use ureteroscopes, are likely to reduce cross-contamination and lower the costs of equipment maintenance, particularly in high-volume centers (Ventimiglia *et al.*, 2022)^[18].

Looking ahead, future innovations are expected to focus on improving the durability of ureteroscopes, optimizing laser lithotripsy techniques, and refining pressure management systems. These advancements will contribute to higher stone-free rates and fewer complications, necessitating regular updates to the guidelines.

4. Limitations of the guidelines

The guidelines are based on a review of current literature and expert consensus, which may not fully capture the variability in patient populations and clinical settings. For example, certain recommendations, such as the use of advanced digital ureteroscopes, may not be feasible in resource-limited settings. Additionally, while the guidelines address common complications, further research is needed to provide detailed recommendations for managing rare or complex cases, such as patients with recurrent stones or significant comorbidities.

The Rapti Academy of Health Sciences guidelines for RIRS provide a valuable framework for standardizing the procedure and ensuring optimal patient outcomes. By following these guidelines, urologists can improve stone-free rates, reduce complications, and enhance overall patient care. As technological advancements continue to evolve, these guidelines will require periodic updates to reflect the latest evidence and best practices.

Conclusion

The comprehensive guidelines for Retrograde Intra Renal Surgery (RIRS) developed by the Rapti Academy of Health Sciences provide a robust framework for the safe and effective management of renal stones. RIRS has proven to be an invaluable minimally invasive procedure, offering high stone-free rates, particularly for renal stones smaller than 2 cm. By following the guidelines that cover patient selection, preoperative evaluation, surgical techniques, and postoperative care, urologists can ensure better outcomes and reduced complications for patients undergoing this procedure.

Including detailed protocols for managing complications and emphasizing the steep learning curve underscores the importance of experience and careful technique in achieving the best possible results. The development of these guidelines also reflects the continuous advancements in technology, particularly in digital ureteroscopes and laser lithotripsy, which are instrumental in improving the safety and efficacy of RIRS.

Despite the effectiveness of RIRS, it remains a technically demanding procedure, requiring extensive training and experience. The guidelines recommend a structured training

approach to ensure that surgeons can achieve proficiency and minimize risks. Additionally, the guidelines acknowledge the importance of staying current with technological advancements, as future innovations will further enhance procedural success.

In conclusion, the guidelines provide a comprehensive and evidence-based approach to performing RIRS. By adhering to these standardized practices, urologists can optimize patient outcomes, minimize complications, and contribute to the wider adoption of RIRS as a standard procedure for managing renal stones. As technology continues to evolve, these guidelines will require regular updates to remain aligned with the latest evidence and innovations in the field.

References

1. Amorim LGCRD, Campos MEC, Dumont LSA, Peñafiel JAR, Abreu ESD, Marchini GS, *et al.* Retrograde intrarenal surgery with or without ureteral access sheath: a systematic review and meta-analysis of randomized controlled trials. *Int Braz J Urol*,2024;50(6):670-682.
2. Asutay MK, Lattarulo M, Liourdi D, Al-Aown AM, Pagonis K, Nedal N, *et al.* Does ureteral access sheath have an impact on ureteral injury? *Urol Ann*,2022;14(1):1-7.
3. Bıçaklıoğlu F, Koparal MURAT, Bulut ENDER. The use of flexible ureteroscopy is required to improve stone-free rates in ureteroscopy for proximal ureteral stones. *Arch Urol Res*, 2021, 5(1).
4. Cooper JL, François N, Sourial MW, Miyagi H, Rose JR, Shields J, *et al.* The impact of ureteral access sheath use on the development of abnormal postoperative upper tract imaging after ureteroscopy. *J Urol*,2020;204(5):976-981.
5. Dasgupta R, Ong TA, Lim J, Rajandram R, Gao X, Hakim L, *et al.* A global perspective of stenting after ureteroscopy: an observational multicenter cohort study. *SIU J*,2021;2(2):96-105.
6. Denstedt J, Basulto-Martínez M, González-Cuenca E. Ureteroscopy and related instruments. In: *The Ureter: A Comprehensive Review*. Cham: Springer International Publishing, 2024, 493-515.
7. Fankhauser CD, Weber D, Müntener M, Poyet C, Sulser T, Hermanns T. Effectiveness of flexible ureterorenoscopy versus extracorporeal shock wave lithotripsy for renal calculi of 5–15 mm: results of a randomized controlled trial. *Eur Urol Open Sci*,2021;25:5-10.
8. Feng D, Zeng X, Han P, Wei X. Comparison of intrarenal pelvic pressure and postoperative fever between standard-and mini-tract percutaneous nephrolithotomy: a systematic review and meta-analysis of randomized controlled trials. *Transl Androl Urol*,2020;9(3):1159.
9. Geraghty RM, Davis NF, Tzelvels L, Lombardo R, Yuan C, Thomas K, *et al.* Best practice in interventional management of urolithiasis: an update from the European Association of Urology Guidelines Panel for Urolithiasis 2022. *Eur Urol Focus*,2023;9(1):199-208.
10. Juliebø-Jones P, Keller EX, Haugland JN, Æsøy MS, Beisland C, Somani BK, Ulvik Ø. Advances in ureteroscopy: new technologies and current innovations in the era of Tailored Endourological Stone Treatment (TEST). *J Clin Urol*,2023;16(3):190-198.

11. Kim CH, Chung DY, Rha KH, Lee JY, Lee SH. Effectiveness of percutaneous nephrolithotomy, retrograde intrarenal surgery, and extracorporeal shock wave lithotripsy for treatment of renal stones: a systematic review and meta-analysis. *Medicina*,2020;57(1):26.
12. Kim HJ, Daignault-Newton S, DiBianco JM, Conrado B, Jafri SM, Seifman B, *et al.* Real-world practice stone-free rates after ureteroscopy: variation and outcomes in a surgical collaborative. *Eur Urol Focus*,2023;9(5):773-780.
13. Lang J, Narendrula A, El-Zawahry A, Sindhwani P, Ekwenna O. Global trends in incidence and burden of urolithiasis from 1990 to 2019: an analysis of global burden of disease study data. *Eur Urol Open Sci*,2022;35:37-46.
14. Lima A, Reeves T, Geraghty R, Pietropaolo A, Whitehurst L, Somani BK. Impact of ureteral access sheath on renal stone treatment: prospective comparative non-randomised outcomes over a 7-year period. *World J Urol*,2020;38:1329-1333.
15. Lildal SK, Osther P, Jung H. Irrigation mechanisms and intrarenal pressure in flexible ureteroscopy. In: *Flexible Uteroscopy*. Singapore: Springer Nature Singapore, 2022, 99-115.
16. Şeçkiner İ, Yılmaz AE, Öztürk M. Fiberoptic or digital? Comparison of the use of two different flexible ureteroscopes in kidney stone treatment. *Endoüroloji Bülteni*,2020;12(2):129-134.
17. Uzun E, Arabaci HB, Ceviz K, Koudonas A, Germiyanoglu RC, Senel S. Development of a new scoring system predicting medical expulsive therapy success on 4–10 mm distal ureteral stones: medical expulsive therapy stone score (METSS). *Urolithiasis*,2023;52(1):8.
18. Ventimiglia E, Smyth N, Doizi S, Jimenez Godinez A, Barghouthy Y, Corrales Acosta MA, *et al.* Can the introduction of single-use flexible ureteroscopes increase the longevity of reusable flexible ureteroscopes at a high volume centre? *World J Urol*,2022;40(1):251-256.
19. Wu G, Sun F, Sun K, Zhang D, Yao H, Wu J, *et al.* Impact of differential ureteral stent diameters on clinical outcomes after ureteroscopy intracorporeal lithotripsy: a systematic review and meta-analysis. *Int J Urol*,2021;28(10):992-999.